

Regioselective Reduction of 3-Methoxymaleimides: an Efficient Method for the Synthesis of Methyl 5- Hydroxytetramates


*Fatihah Issa, Joshua Fischer, Peter Turner and Mark J. Coster**

School of Chemistry, The University of Sydney, NSW 2006, Australia

m.coster@chem.usyd.edu.au



RECEIVED DATE (to be automatically inserted after your manuscript is accepted if required according to the journal that you are submitting your paper to)

Regioselective Reduction of 3-Methoxymaleimides



Abstract. 3-Methoxymaleimide and various *N*-alkyl-3-methoxymaleimides, synthesized by base-promoted *N*-alkylation of 3-methoxymaleimide, were reduced using sodium borohydride with complete regioselectivity. The resultant methyl 5-hydroxytetramates are useful intermediates in the synthesis of a variety of tetramate derivatives.


Tetramic acids (pyrrolidine-2,4-diones) **1** are a common structural feature in a number of biologically-active natural products, including compounds with antibacterial, antifungal and cytotoxic properties.¹ 5-Hydroxytetramic acids (5-hydroxypyrrolidine-2,4-diones) **2**, which exhibit an *N*-acylhemiaminal functionality, are less common, though still represented in the HIV-integrase inhibitory and immunomodulatory/antibacterial natural products integramycin (**3**)² and delaminomycin A (**4**),³ respectively. A direct and convenient approach to the synthesis of 5-hydroxytetramic acids would rely on the regioselective reduction of a suitable maleimide derivative. Given the ready availability of 3-methoxymaleimide (**5a**),⁴ and the known facility with which *O*-alkyl tetramates can be hydrolyzed to tetramic acids,⁵ the regioselective reduction of 3-methoxymaleimides and subsequent hydrolysis appeared to be a potentially convenient route to 5-hydroxytetramic acids.



3-Methoxymaleimide (**5a**) is readily available from maleimide by a convenient two step procedure.⁴ A variety of *N*-alkyl-3-methoxymaleimides **5b-f** were prepared by base-promoted

alkylation of **5a** (Table 1). The potassium carbonate promoted *N*-benzylation⁶ and *N*-methylation^{7,8} of maleimides with benzyl bromide and iodomethane, respectively, in acetone solvent, have previously been reported. Reaction of **5a** with a variety of alkylating agents under these conditions (Method A), provided the desired *N*-alkyl-3-methoxymaleimides **5b-f** in good yields (87–92%) when MeI, EtI or PMBCl were employed (entries 1, 2 and 5). The use of MeCN as solvent markedly improved the yield of **5e** (99% vs. 55% with acetone), however this procedure with either solvent did not prove satisfactory for the allylation of **5a** to give **5d** (10% with acetone vs. 2% with MeCN). An alternative procedure, employing cesium fluoride in DMF (Method B), inspired by the facile CsF-promoted *N*-alkylation of phthalimide with alkyl chlorides reported by Clark and Miller,⁹ proved to be facile for most alkylations, including the reaction of **5a** with allyl chloride (84%). 3-Methoxy-*N*-phenylmaleimide (**5g**) was prepared in three steps from *N*-phenylmaleimide by the method of Argade,¹⁰ as another substrate for the regioselective reduction.

Table 1. Alkylation of methoxymaleimide (**5a**)^a


				
entry	R–X	product	Method A yield (%) ^b	Method B yield (%) ^b
1	MeI	5b	87	90
2	EtI	5c	92	61
3	Allyl chloride	5d	10	84

4	BnCl	5e	99 ^c	94
5	PMBCl	5f	91	68

^a **Method A:** **5a**, RX (2–2.9 equiv.), K₂CO₃ (1.2 equiv.), acetone, reflux, 17–90 h; **Method B:** **5a**, RX (3 equiv.), CsF (3 equiv.), DMF, 30–38 °C, 24 h. ^b Isolated yields. ^c MeCN as solvent, 50 °C (55% yield with acetone).

Citraconimide (3-methylmaleimide), and *N*-alkyl derivatives, have previously been shown to undergo regioselective reduction with sodium borohydride to give predominantly the products of attack at the more hindered C2 position (88 : 12 → >99 : <1), whereas a reversal in regioselectivity was observed with NaBH₄ in the presence of cerium(III) chloride heptahydrate (CeCl₃·7H₂O) or by using diisobutylaluminum hydride (DIBAL-H).¹¹ We reasoned that delocalisation of a lone pair on the methoxy oxygen of **5a-g** would significantly decrease the reactivity of the C5 carbonyl group towards nucleophilic attack by hydride reducing reagents, and that for this reason, the reduction of these systems should be particularly regioselective.¹² A survey of common reducing agents showed DIBAL-H, in THF or CH₂Cl₂, to be surprisingly non-selective in its reaction with **5e**, providing *ca.* 1:1 mixtures of regioisomeric reduction products. Reduction of **5e** with LiAlH₄ gave an 88:12 mixture of products resulting from attack at C2 and C5 respectively, and appreciable quantities of over-reduced compounds. The use of NaBH₄–CeCl₃ improved the regioselectivity to 97:3 in favour of the C2 reduced compound, however, the use of NaBH₄ by itself, in THF–H₂O at 0 °C proved particularly effective and convenient, providing **6e** as the sole isolated regioisomer in 83% yield (Table 2, entry 5). In all cases studied (Table 2), these conditions provided good yields (70–87%) of the methyl 5-hydroxytetramates **6a-g**, with no evidence for the formation of the regioisomeric products, resulting from borohydride attack at C5 of **5a-g**.

Table 2. Regioselective reduction of **5a-g**



entry	substrate	R	t/min	product ^a
1	5a	H	30	6a (87%)
2	5b	Me	45	6b (87%)
3	5c	Et	90	6c (84%)
4	5d	allyl	30	6d (85%)
5	5e	Bn	100	6e (83%)
6	5f	PMB	180	6f (76%)
7	5g	Ph	60	6g (70%)


^a Isolated yields.

The physical and spectroscopic data, *eg.* ¹H and ¹³C NMR, for **6a** and **6b** match that reported previously,^{12,13} while the structure of **6e** was confirmed by single crystal X-ray diffraction, the ORTEP depiction is provided in the Supporting Information. The structures of **6c-d,f-g**, and therefore the regioselectivity of the reduction, were assigned by analogy with the aforementioned, closely related examples.¹⁴

Attempts to hydrolyze methyl 5-hydroxytetramates **6a** and **6e** (aq. HCl) have thus far been thwarted by difficulties with purification and/or instability of the resultant 5-hydroxytetramic acids. It is possible that 3-acyl derivatives, such as the natural products integramycin (**3**) and delaminomycin (**4**), are inherently more stable, and as such, future work will be directed towards these types of systems. Nevertheless, methyl 5-hydroxytetramates **6a-g** afforded by the regioselective reduction reported herein, are versatile building blocks for the synthesis of a range of tetramate derivatives (Scheme 1). Treatment of **6a** and **6e** with methanolic HCl afforded

methyl 5-methoxytetramates **7** and **8**, in 100% and 89% yields, respectively. Reduction of **6e** with Et_3SiH and $\text{BF}_3 \cdot \text{OEt}_2$ to afford **9** (96% yield), demonstrates that access to methyl *N*-alkyltetramates can be achieved. Acetylation of **6e** (Ac_2O , pyridine, DMAP) followed by treatment of the resultant acetate with allyltrimethylsilane and $\text{BF}_3 \cdot \text{OEt}_2$ provided 5-allyltetramate **10** in good yield (80% over two steps).

Scheme 1. Synthesis of Methyl Tetramate Derivatives



Experimental Section

Method A representative procedure: To a solution of 3-methoxymaleimide (**5a**) (40.0 mg, 0.315 mmol) in MeCN (0.52 mL) was added K₂CO₃ (52.2 mg, 0.378 mmol, 1.2 equiv.) and then BnCl (72 μ L, 0.63 mmol, 2.0 equiv.) dropwise. The mixture was heated to 50 °C until t.l.c analysis indicated complete conversion of the starting material (90 h). After cooling to rt the reaction mixture was concentrated *in vacuo*, then partitioned between half-saturated aq. NH₄Cl and EtOAc. The layers were separated and the aqueous phase was extracted with EtOAc (2 \times). The combined organic layers were dried (MgSO₄), filtered, then the filtrate was concentrated *in vacuo*. Purification of the residue by flash chromatography (hexane/EtOAc 60:40) gave 1-benzyl-3-methoxy-1*H*-pyrrole-2,5-dione (**5e**) as a colourless solid (67.6 mg, 99%): mp 68-69 °C; IR (thin film) 3117, 3032, 1709, 1639 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.27 (m, 5H), 5.39 (s, 1H), 4.62 (s, 2H), 3.85 (s, 3H); ¹³C NMR (75.5 MHz, CDCl₃) δ 169.6 (C), 165.0 (C), 160.8 (C), 136.1 (C), 128.4 (CH), 128.0 (CH), 127.5 (CH), 96.2 (CH), 58.8 (CH₃), 40.9 (CH₂); MS (ESI) *m/z* 240 [M + Na]⁺, 218 [M + H]⁺; HRMS (EI) calcd for C₁₂H₁₁NO₃ (M⁺) 217.0739, found 217.0737.

Method B representative procedure: To a mixture of 3-methoxymaleimide (**5a**) (63.5 mg, 0.500 mmol) and anhydrous CsF (228 mg, 1.50 mmol, 3.0 equiv.) was added DMF (1.67 mL), followed by allyl chloride (122 μ L, 1.50 mmol, 3.0 equiv.). The mixture was heated at 30-38 °C for 24 h. The suspension was diluted with half-saturated aq. NaHCO₃ solution and EtOAc, then the layers were separated. The aqueous phase was extracted with EtOAc (5 \times) and the combined organic extracts were dried (MgSO₄), filtered and concentrated *in vacuo*. Purification of the residue by flash chromatography (hexane/EtOAc 60:40) provided 1-allyl-3-methoxy-1*H*-pyrrole-2,5-dione (**5d**) as a colourless solid (70.0 mg, 84%): mp 61-63 °C; IR (thin film) 3111, 1705, 1626 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 5.80 (ddt, *J* = 17.2, 15.7, 5.5 Hz, 1H), 5.43 (s, 1H),

5.21 (dq, $J = 7.8, 1.5$ Hz, 1H), 5.14 (m, 1H), 4.11 (dt, $J = 5.6, 1.5$ Hz, 2H), 3.94 (s, 3H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 169.6 (C), 165.1 (C), 160.9 (C), 131.5 (CH), 117.5 (CH_2), 96.2 (CH), 58.9 (CH_3), 39.6 (CH_2); MS (ESI) m/z 190 $[\text{M} + \text{Na}]^+$, 168 $[\text{M} + \text{H}]^+$; HRMS (ESI) calcd for $\text{C}_8\text{H}_9\text{NO}_3\text{Na}$ $[\text{M} + \text{Na}]^+$ 190.0475, found 190.0477.

Representative procedure for the regioselective reduction using NaBH_4 : To a cooled (0 °C) solution of **5e** (65.5 mg, 0.302 mmol) in 2:1 THF (0.3 mL) and H_2O (0.15 mL), was added NaBH_4 (12 mg, 0.31 mmol, 1.0 equiv.). The resulting solution was stirred at 0 °C for 100 min, then quenched by the dropwise addition of acetone (1 mL). Silica gel (0.7 g) was added and the volatile components were removed *in vacuo*. Purification by flash chromatography (MeOH/ CH_2Cl_2 5:95) afforded 1-benzyl-5-hydroxy-4-methoxy-1*H*-pyrrol-2(5*H*)-one (**6e**) as a colourless solid (54.9 mg, 83%): mp 134-138 °C; IR (thin film) 3130, 3032, 1655, 1643 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.30 (m, 5H), 5.06 (br d, $J = \sim 9$ Hz, 1H), 5.00 (s, 1H), 4.94 (d, $J = 15.1$ Hz, 1H), 4.17 (d, $J = 15.1$ Hz, 1H), 3.80 (s, 3H), 3.68 (br d, $J = \sim 9$ Hz, 1H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 173.9 (C), 170.4 (C), 137.2 (C), 128.7 (CH), 128.2 (CH), 127.5 (CH), 93.2 (CH), 80.0 (CH), 58.4 (CH_3), 42.3 (CH_2); MS (ESI) m/z 242 $[\text{M} + \text{Na}]^+$, 220 $[\text{M} + \text{H}]^+$, 202 $[\text{M} - \text{OH}]^+$; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{13}\text{NO}_3\text{Na}$ $[\text{M} + \text{Na}]^+$ 242.0788, found 242.0784.

1-Benzyl-4,5-dimethoxy-1*H*-pyrrol-2(5*H*)-one (8**).** To a cooled (0 °C) solution of **6e** (100 mg, 0.456 mmol) in MeOH (4.6 mL) was added TMSCl (174 μL , 1.37 mmol, 3.0 equiv.). The reaction was warmed to rt and allowed to stir for 21 h, then the mixture was concentrated *in vacuo*. The residue was dissolved in EtOAc and the organic phase was washed with half-saturated aq. NaHCO_3 solution, then brine. The organic phase was dried (Na_2SO_4), filtered and concentrated. Purification of the residue by flash chromatography (EtOAc/acetone 95:5) afforded **8** as a colourless oil (94.8 mg, 89%): IR (thin film) 3107, 3028, 1703, 1634 cm^{-1} ; ^1H NMR (300

MHz, CDCl₃) δ 7.34-7.24 (m, 5H), 5.14 (s, 1H), 5.04 (s, 1H), 4.96 (d, J = 14.9 Hz, 1H), 4.02 (d, J = 14.9 Hz, 1H), 3.80 (s, 3H), 3.11 (s, 3H); ¹³C NMR (75.5 MHz, CDCl₃) δ 171.4 (C), 170.0 (C), 137.1 (C), 128.5 (CH), 128.3 (CH), 127.4 (CH), 95.2 (CH), 84.9 (CH), 58.2 (CH₃), 50.4 (CH₃), 42.5 (CH₂); MS (ESI) m/z 234 [M + H]⁺; HRMS (ESI) calcd for C₁₃H₁₆NO₃ [M + H]⁺ 234.1125, found 234.1131.

1-Benzyl-4-methoxy-1H-pyrrol-2(5H)-one (9). To a cooled (0 °C) suspension of **6e** (40.5 mg, 0.185 mmol, 1.0 equiv.) in CH₂Cl₂ (1.3 mL) was added triethylsilane (59 μ L, 0.37 mmol, 2.0 equiv.), followed by BF₃·OEt₂ (47.0 μ L, 0.370 mmol, 2.0 equiv.). The mixture was slowly warmed to rt and stirred for 24 h. The solution was quenched with saturated aq. K₂CO₃ solution, then the layers were separated. The aqueous layer was extracted with CH₂Cl₂ (3 \times) and the combined organic portions were dried (Na₂SO₄), filtered and concentrated. Purification by flash chromatography (EtOAc/acetone 95:5) afforded **9** as a colourless solid (36.1 mg, 96%): mp 44-45 °C; IR (thin film) 3028, 1674, 1624 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.34-7.22 (m, 5H), 5.10 (s, 1H), 4.57 (s, 2H), 3.76 (s, 3H), 3.72 (s, 2H); ¹³C NMR (75.5 MHz, CDCl₃) δ 173.3 (C), 172.0 (C), 137.4 (C), 128.6 (CH), 127.8 (CH), 127.4 (CH), 94.1 (CH), 58.0 (CH₃), 49.8 (CH₂), 45.3 (CH₂); MS (ESI) m/z 226 [M + Na]⁺, 204 [M + H]⁺; HRMS (ESI) calcd for C₁₂H₁₃NO₂Na [M + Na]⁺ 226.0839, found 226.0836.

5-Allyl-1-benzyl-4-methoxy-1H-pyrrol-2(5H)-one (10). To a solution of **6e** (76.3 mg, 0.348 mmol, 1.0 equiv.) in pyridine (7.0 mL) at rt was added Ac₂O (329 μ L, 3.48 mmol, 10 equiv.) followed by DMAP (4.3 mg, 0.0348 mmol, 0.1 equiv.). After stirring for 3 h, the reaction was quenched with water and extracted with Et₂O (5 \times). The combined organic layers were dried (Na₂SO₄), filtered and concentrated. Purification by flash chromatography (hexane/EtOAc 60:40) afforded 5-acetoxy-1-benzyl-4-methoxy-1H-pyrrol-2(5H)-one as a colourless solid (77.5

mg, 85%): mp 102-104 °C; IR (thin film) 3115, 1749, 1711, 1641 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.30-7.23 (m, 5H), 6.42 (s, 1H), 5.13 (s, 1H), 4.59 (d, *J* = 15.3 Hz, 1H), 4.41 (d, *J* = 15.3 Hz, 1H), 3.80 (s, 3H), 1.88 (s, 3H); ¹³C NMR (75.5 MHz, CDCl₃) δ 171.2 (C), 170.5 (C), 170.2 (C), 137.2 (C), 128.4 (CH), 128.0 (CH), 127.3 (CH), 94.5 (CH), 79.2 (CH), 58.5 (CH₃), 43.8 (CH₂), 20.4 (CH₃); MS (ESI) *m/z* 284 [M + Na]⁺; HRMS (ESI) calcd for C₁₄H₁₅NO₄Na [M + Na]⁺ 284.0894, found 284.0897. To a cooled (0 °C) solution of 5-acetoxy-1-benzyl-4-methoxy-1*H*-pyrrol-2(5*H*)-one (20.5 mg, 0.0785 mmol, 1.0 equiv.), from the above procedure, in CH₂Cl₂ (0.8 mL), was added BF₃·OEt₂ (39.8 μL, 0.314 mmol, 4.0 equiv.). After stirring at 0 °C for 10 min, allyltrimethylsilane (37.4 μL, 0.236 mmol, 3.0 equiv.) was added dropwise via syringe. The reaction was stirred at 0 °C for 7 h, then warmed to rt and stirred for another 16 h. After pouring the mixture into water (1 mL), the layers were separated and the aqueous phase was extracted with CH₂Cl₂ (3 ×). The combined organic portions were washed with brine, dried (Na₂SO₄), filtered and concentrated *in vacuo*. Purification by flash chromatography (hexane/acetone 60:40) afforded **10** as a colourless oil (17.9 mg, 94%): IR (thin film) 3076, 3028, 1680, 1626 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 7.34-7.21 (m, 5H), 5.51 (ddt, *J* = 17.1, 10.2, 7.1 Hz, 1H), 5.17 (d, *J* = 15.4 Hz, 1H), 5.10 (s, 1H), 5.07 (m, 1H), 5.06 (m, 1H), 3.98 (d, *J* = 15.4 Hz, 1H), 3.87 (t, *J* = 4.3 Hz, 1H), 3.76 (s, 3H), 2.57-2.38 (m, 2H); ¹³C NMR (75.5 MHz, CDCl₃) δ 175.3 (C), 171.7 (C), 137.6 (C), 131.0 (CH), 128.6 (CH), 127.9 (CH), 127.4 (CH), 118.8 (CH₂), 94.2 (CH), 58.5 (CH₃), 58.0 (CH), 43.2 (CH₂), 32.6 (CH₂); MS (ESI) *m/z* 266 [M + Na]⁺, 244 [M + H]⁺; HRMS (ESI) calcd for C₁₅H₁₇NO₂Na [M + Na]⁺ 266.1152, found 266.1152.

Acknowledgment. We thank the Australian Research Council for funding, and CSIRO for a scholarship to JF.

Supporting Information Available: Experimental procedures and characterization data for new compounds not included in the Experimental Section; ORTEP depiction and X-ray crystallographic data (CIF) for **6e**; ^1H and ^{13}C NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Royles, B. J. L. *Chem. Rev.* **1995**, *95*, 1981-2001.
- (2) Singh, S. B.; Zink, D. L.; Heimbach, B.; Genilloud, O.; Teran, A.; Silverman, K. C.; Lingham, R. B.; Felock, P.; Hazuda, D. J. *Org. Lett.* **2002**, *4*, 1123-1126.
- (3) (a) Ishizuka, M.; Kawatsu, M.; Yamashita, T.; Ueno, M.; Takeuchi, T. *Int. J. Immunopharmacol.* **1995**, *17*, 133-139; (b) Ueno, M.; Amemiya, M.; Iijima, M.; Osono, M.; Masuda, T.; Kinoshita, N.; Ikeda, T.; Iinuma, H.; Hamada, M.; Ishizuka, M.; Takeuchi, T. *J. Antibiot.* **1993**, *46*, 719-727; (c) Ueno, M.; Someno, T.; Sawa, R.; Iinuma, H.; Naganawa, H.; Ishizuka, M.; Takeuchi, T. *J. Antibiot.* **1993**, *46*, 979-984; (d) Ueno, M.; Someno, T.; Sawa, R.; Iinuma, H.; Naganawa, H.; Ishizuka, M.; Takeuchi, T. *J. Antibiot.* **1993**, *46*, 1020-1023; (e) Ueno, M.; Amemiya, M.; Yamazaki, K.; Iijima, M.; Osono, M.; Someno, T.; Iinuma, H.; Hamada, M.; Ishizuka, M.; Takeuchi, T. *J. Antibiot.* **1993**, *46*, 1156-1162; (f) Ueno, M.; Yoshinaga, I.; Amemiya, M.; Someno, T.; Iinuma, H.; Ishizuaki, M.; Takeuchi, T. *J. Antibiot.* **1993**, *46*, 1390-1396.
- (4) Booker-Milburn, K. I.; Baker, J. R.; Bruce, I. *Org. Lett.* **2004**, *6*, 1481-1484.
- (5) Baenziger, M.; McGarrity, J. F.; Meul, T. *J. Org. Chem.* **1993**, *58*, 4010-4012.
- (6) Joyce, R. P.; Gainor, J. A.; Weinreb, S. M. *J. Org. Chem.* **1987**, *52*, 1177-1185.

(7) Lakatosh, S. A.; Balzarini, J.; Andrei, G.; Snoeck, R.; De Clercq, E.; Preobrazhenskaya, M. N. *J. Antibiot.* **2002**, *55*, 768-773.

(8) (a) Tominaga, Y.; Shigemitsu, Y.; Sasaki, K. *J. Heterocycl. Chem.* **2002**, *39*, 571-591; (b) Shigemitsu, Y.; Tominaga, Y. *Heterocycles* **2001**, *55*, 2257-2260.

(9) Clark, J. H.; Miller, J. M. *J. Am. Chem. Soc.* **1977**, *99*, 498-504.

(10) Sahoo, M. K.; Mhaske, S. B.; Argade, N. P. *Synthesis* **2003**, 346-349.

(11) Mase, N.; Nishi, T.; Hiyoshi, M.; Ichihara, K.; Bessho, J.; Yoda, H.; Takabe, K. *J. Chem. Soc., Perkin Trans. 1* **2002**, 707-709.

(12) Stachel, H.-D.; Schachtner, J.; Seidel, J. *Z. Naturforsch., B: Chem. Sci.* **1996**, *51*, 409-416.

(13) Farina, F.; Victoria Martin, M.; Carmen Paredes, M.; Carmen Ortega, M.; Tito, A. *Heterocycles* **1984**, *22*, 1733-1739.

(14) The chemical shift of the olefinic methine carbon in the two possible regioisomers is also diagnostic, δ_C 93.2 for **6e**, in which this carbon is α - to the carbonyl, *cf.* δ_C 106.8 for the other regioisomer, in which this carbon is β - to the carbonyl. See the Supporting Information for spectroscopic data for this regioisomeric product.