

The design, synthesis, and anti-inflammatory evaluation of a drug-like library based on the natural product valerenic acid

Folake A. Egbewande,^a Niclas Nilsson,^b Jonathan M. White,^c Mark J. Coster,^a and Rohan A. Davis^{*,a}

^aGriffith Institute for Drug Discovery, Griffith University, Brisbane, QLD 4111, Australia.

^bLEO Pharma A/S, Industriparken 55, 2750 Ballerup, Denmark.

^cSchool of Chemistry and Bio21 Institute, The University of Melbourne, VIC 3010, Australia.

Corresponding Author

* Tel: +61-7-3735-6043. Fax: +61-7-3735-6001. E-mail: r.davis@griffith.edu.au

ABSTRACT: The plant natural product, valerenic acid (**1**) was chosen as a desirable scaffold for the generation of a novel screening library due to its drug-like physicochemical parameters

(such as log P, hydrogen bond donor/acceptor counts, and molecular weight). An 11-membered amide library (**2–12**) was subsequently generated using parallel solution-phase synthesis. The chemical structures of all semi-synthetic analogues (**2–12**) were elucidated following analysis of the NMR, MS, UV and IR data. The structures of compounds **8** and **11** were also confirmed by X-ray crystallographic analysis. All library members were evaluated for their ability to inhibit the release of IL-8 and TNF- α . Six analogues showed moderate activity in the IL-8 assay with IC₅₀ values of 2.8–8.3 μ M, while none of the tested compounds showed any significant effect on inhibiting TNF- α release.

Nature has played an important role in drug discovery and development,¹ yet it still remains an under-investigated source of unique chemical scaffolds for semi-synthetic library generation, and subsequent bioassay screening for hit or lead molecules.^{2, 3} Natural products

(NPs) are a pre-validated source of small molecules that can be used for the design of unique biologically active compounds, because they have been optimized via natural evolution for maximum interactions with biosynthetic enzymes.^{2,4,5} For many decades, NPs have impacted pharmaceutical research and development by providing unique chemical diversity and complexity that was exploited as either NP drugs or as starting materials (i.e. hits/leads).^{6,7} Between 1981 and 2014, 387 NP-based drugs were approved for use worldwide, 67 of which were unaltered NP, while 320 were NP-derived drugs.¹

The genus *Valeriana* is made up of about 200 species, some of which are endemic to Europe and Asia, others to North America and South America.^{8,9} In many cultures, the roots and rhizomes of various species are used traditionally for the treatment of insomnia and anxiety.¹⁰ This genus has been shown to display a wide range of biological effects including, anti-HIV,¹¹ cytotoxicity,¹² anti-convulsant,¹³ and anti-hypertensive activities.¹⁴ Previous investigations of this genus showed the presence of iridoids, sesquiterpenoids, flavone glycosides, lignans, and alkaloids.^{15, 16, 17, 18, 19, 20}

Valeriana officinalis is a well-known source of the bioactive sesquiterpene, valerenic acid (**1**). Pharmacological studies have shown that valerian extracts allosterically modulate GABA_A receptors, with valerenic acid (**1**) described as one of the active principles underlying this observed effect.²¹ Several investigations have revealed various biological activities observed for valerenic acid and its derivatives, namely cytotoxicity,¹² anxiolytic,^{13, 22, 23, 24} and anti-inflammatory activities.²⁵ *V. officinalis* has been widely researched, with the aim of understanding the activity, which has been observed *in vivo* and *in vitro*. Yet, this plant and its chemistry are still subjects of considerable investigation, aimed at finding new biological targets and effects e.g. anticoronaryspastic and antibronchospastic activities.¹⁴

Our research focuses on the design and semi-synthesis of drug discovery libraries based on unique NP scaffolds from various biota sources, such as fungi, plants, and marine

invertebrates.^{26, 27, 28, 29, 30} The ultimate goal of such libraries is to assist in the identification of hit or lead compounds that impact the NP drug discovery process. The use of NP scaffolds as the starting point for focused library synthesis is proving to be a powerful tool for NP-based drug discovery.³¹

As part of our continuing efforts to contribute to knowledge in this area of research, valerenic acid (**1**) was chosen for library generation and medicinal chemistry studies. Our strategy involved the use of commercially available valerenic acid (**1**), which has the advantages of saving both time and cost, as it bypasses the *de novo* synthesis for scaffold production. This sesquiterpenoid was an attractive NP scaffold for synthetic studies since it was commercially available, has a low molecular weight (MW) (234 Da), multiple stereogenic centers (n = 3), favourable calculated log P (cLogP, 3.21), and a functional group (carboxylic acid) that is suitable for chemical modification.

Herein we report the design, parallel solution-phase synthesis and the anti-inflammatory evaluation of a library of semi-synthetic derivatives based on the NP scaffold, valerenic acid (**1**).

The amide functional group plays a major role in the composition of biological systems.³² Being the essential chemical bond found in peptides and proteins,³³ this moiety is ubiquitous in life, as proteins are significant in many biological processes such as enzymatic catalysis, transport/storage, immune protection, and mechanical support.³² This functionality also plays a key role in medicinal chemistry, widely occurring in biologically active compounds and pharmaceuticals, such as the antifungal drug anidulafungin,³⁴ the anticancer drugs flutamide and bicalutamide,¹ the multikinase inhibitor and antitumor agent sorafenib,² and the antibiotic tigecycline.³⁴

Prior to commencing synthesis on scaffold **1**, 26 commercially available amines were initially selected, and a virtual analogue (VA) library generated (VA1–VA26), which was

subsequently analyzed using ChemDraw Ultra (see Supplementary material S66 for chemical structures of VA1–VA26).³⁵ Physicochemical parameters such as cLog P, hydrogen bond donors (HBD), hydrogen bond acceptors (HBA), and molecular weight (MW) of these VAs were determined *in silico*. The molecules with desirable physicochemical properties, as described by Lipinski’s “Rule of Five” for orally bioavailable drug-like compounds (HBD \leq 5, HBA \leq 10, MW \leq 500 and Log P \leq 5),³⁶ were subsequently chosen for synthesis (see Supplementary material Table S65). VA1–VA11 were prioritized for synthesis, since they all had minimal or no “Rule of Five” violations.

Literature reports have shown the usage and advantages of coupling agents in amide bond formation.^{32, 33, 37, 38} However, making an appropriate choice of coupling reagent is often demanding, as a result of the plethora of reagents already reported.^{32, 33} The phenethylamine derivative (**2**) was chosen as the first synthetic target for coupling optimization studies. Trial reactions were initially attempted using the commercially available valerenic acid and three coupling reagents which included, *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide (EDCI),²⁶ *N,N,N',N'*-tetramethylchloroformamidinium hexafluorophosphate (TCFH),³⁹ and 1-chloro-*N,N,2*-trimethyl-1-propenyl-amine (Ghosez’s reagent),⁴⁰ with yields of 6%, 12%, and 91% respectively. Due to the superior yield associated with Ghosez’s reagent, this was chosen as the coupling reagent of choice for all other amidation reactions undertaken on scaffold **1**.

Treatment of NP scaffold **1** with 11 primary amines, and Ghosez’s reagent afforded the secondary amides (**2–12**) (Fig. 1 and Scheme 1) in moderate to excellent yields (32–99%). The structures of all the amide analogues were determined following 1D/2D NMR and (+)-HRESIMS data analysis (see Supplementary material). While the synthesis of valerenic acid amide analogues have been previously reported by other researchers,^{40, 41} this is the first reported synthesis and characterization of amides **2–12**.

An example of the NMR characterization of compound **2** is given below. Briefly, the ^1H NMR spectrum in $\text{DMSO-}d_6$ indicated the presence of two methylenes [δ_{H} 2.74 (H-17), and 3.30 (H-16)], three aromatic protons [δ_{H} 7.18 (H-21), 7.20 (H-19 and H-23) and 7.28 (H-20 and H-22)], and an amide proton (δ_{H} 7.86); these data were consistent for amidation of the valerenic acid scaffold. Analysis of the COSY spectrum (See Fig. 2 and Supplementary material Fig. S4) of this compound identified three spin systems. Fragments H-1/H₂-2/H₂-3/H-6/H-7/H₂-8/H₂-9/H-10/H₃-14 was the first spin system identified. The second was located between the two methylene protons at H₂-16/H₂-17, and NH, with HMBC correlations from H₂-16 (δ_{H} 3.30) to C-12 (δ_{C} 168.6), C-17 (δ_{C} 35.2), and C-18 (δ_{C} 139.7), and from H₂-17 (δ_{H} 2.74) to C-16 (δ_{C} 40.7), C-18 (δ_{C} 139.7), and C-19/23 (δ_{C} 128.6). The amide proton also showed HMBC correlations to C-16 (δ_{C} 40.7) and C-12 (δ_{C} 168.6). These HMBC correlations in conjunction with the ROESY correlations (Fig. 2) confirmed the formation of the desired product, as well as the conservation of the configuration of the molecule. The last spin system was placed in the aromatic part of the molecule H-19/H-20/H-21/H-22/H-23.

These data enabled the chemical structure of **2** to be unambiguously assigned. The relative configuration of compound **2** was shown to be identical to the NP scaffold following the analysis of the ROESY and ^1H - ^1H coupling constant data, and the magnitude of the ^1H NMR chemical shifts.

Slow evaporation of solutions (*n*-hexane/EtOAc) of **8** and **11** resulted in crystals suitable for X-ray diffraction. The absolute configuration of valerenic acid has been previously determined using circular dichroism,⁴² and this was consistent with the X-ray data of compounds **8** and **11**, which were determined using a Cu source (Fig. 3).

All library members (**2–12**) together with the NP scaffold (**1**) were submitted to LEO Pharma's Open Innovation platform, which is an open collaborative initiative to explore skin inflammation-related science by allowing external partners access to advanced *in vitro*

assays.⁴³ As part of this initiative, the library was tested in two separate anti-inflammatory *in vitro* assays. One assay was looking for inhibitors of lipopolysaccharide (LPS)-induced tumor necrosis factor alpha (TNF- α) release from human primary peripheral blood mononuclear cell (PBMC), while the other assay was searching for small molecules that inhibited interleukin-8 (IL-8) release from IL-17 and TNF- α -induced primary human keratinocytes.

Psoriasis is a chronic, immune-mediated inflammatory skin disorder that occurs in approximately 3% of the world's population,⁴⁴ with 20% of patients suffering from moderate-to-severe disease.^{45, 46} Symptoms include skin lesions, inflammatory plaques and silvery scaling due to keratinocyte hyperproliferation. Key disease mediators include tumor necrosis factor-alpha (TNF- α), and interleukin-17 (IL-17) which induces an inflammatory response measured as an increase in IL-8 secretion.⁴⁷ All compounds were tested for their ability to inhibit this inflammatory response in primary human keratinocytes. In addition, to distinguish a specific inhibition from a cytotoxic effect, a secondary readout, cell viability, was measured to determine possible general cytotoxic mechanisms.

Several compounds (**2–8**) showed moderate ability to inhibit IL-17 and TNF- α induced IL-8 release from primary human keratinocytes, but only at the highest test concentrations. Furthermore, compounds **2–4** exhibited only a very low effect on cell viability, strengthening the relevance of the effect on IL-8 release and indicating a possible scaffold for further development of early compounds for psoriasis treatment. All biological *in vitro* data is presented in Table 1.

In regards to structure-activity relationships, the aliphatic amides, **11** and **12** showed low activity in inhibiting IL-8 release from primary human keratinocytes; substituting with a cyclic side chain such as that in the morpholinoethyl amide derivative, **10** further reduced the potency. However, ~~replacing the side chain with~~introduction of -an aromatic ring into the side chain, such as those found in **2–8**, typically improved the activity. For instance, the benzyl

amide series (**5–8**) showed the greatest inhibition of IL-8 release, but all molecules displayed a very high effect on cell viability, compared to the phenethyl amides **2–4**, which showed low effects.

For example, benzyl amide (**5**) and *ortho*-fluorobenzyl amide (**6**) have similar activities, 2.9 and 2.8 μM , respectively. However, ~~changing moving~~ fluorine from the *ortho*- to the *meta*- (**7**) ~~and or~~ *para*- (**8**) positions decreased the activities by 1.6-fold and 1.7-fold, respectively. ~~On the other hand~~ Interestingly, *para*-fluorophenethyl amide (**3**) was inactive when compared with *para*-fluorobenzyl amide (**8**).

Introducing nitrogen into the aromatic ring [i.e the pyridylethyl amide (**9**)] resulted in a loss of activity. Furthermore, substituting the phenethyl amide unit in **2** with a benzyl amide motif in **5**, resulted in a 2.6-fold increase in IL-8 activity, but a 6.5-fold decrease in cell viability. The same trend was observed when fluorine was introduced into the *para*-positions of **3** and **8**. This indicates that the benzyl amides showed better activity than the phenethyl amides in inhibiting IL-17 and TNF- α induced IL-8 release from primary human keratinocytes; but the later series showed better selectivity.

Table 1. Biological Activity of Compounds **1–1312**

Compound	KC-IL-8		KC-Viability	
	Abs. IC ₅₀ (μM)	Max ^a (%)	Abs.IC ₅₀ (μM)	Max ^a (%)
1	- ^b	4	N/A	15


2	7.5	64	N/A	11
3	N/A	44	N/A	17
4	8.3	60	N/A	24
5	2.9	84	9.0	72
6	2.8	85	8.7	72
7	4.7	79	9.6	67
8	4.9	70	9.6	61
9	N/A	14	N/A	20
10	N/A	17	N/A	12
11	N/A	22	N/A	9
12	N/A	30	N/A	7

^a Max denotes the maximal inhibition of either the IL-8 release or the cell viability, where the higher number represent a more efficacious biological effect. Typically, the Max effect is achieved at the highest test concentration which is 10 μ M.

^b IC₅₀ value was not calculated.

The endotoxin LPS is one of the most potent inducers of inflammatory response. LPS activates the Toll-like Receptor 4 (TLR4), which is responsible for activating the innate immune system, resulting in the release of the pro-inflammatory cytokine TNF- α .⁴⁸ All compounds were tested for the ability to inhibit LPS-induced TNF- α release from primary human peripheral blood mononuclear cell (PBMC), as well as general effect on cell viability. None of the tested compounds showed any significant effect (< 15% inhibition) on reducing the LPS-induced release of TNF- α from primary human PBMC.

In summary, 12 amide derivatives based on the valerenic acid scaffold were synthesised and evaluated for their *in vitro* anti-inflammatory activity in IL-8 and TNF- α assays. Six analogues showed moderate activity in the IL-8 assay with IC₅₀ values of 2.8–8.3 μ M while none of the tested compounds showed any significant effect on TNF- α .



Scheme 1. Reagents and conditions. (i) anhydrous CH_2Cl_2 , 1-Chloro-*N,N*,2-trimethylpropenylamine, $0\text{ }^\circ\text{C} \rightarrow \text{rt}$, 4 h (ii) amine, $0\text{ }^\circ\text{C} \rightarrow \text{rt}$, 16 h (yield = 32–99%).




Fig. 1 Chemical structures of the amide library **2–12**.




Fig. 2 COSY (bold line), key HMBC (→), and ROESY (↔) correlations for compound **2**.




Fig. 3 **A**, ORTEP of **8** showing two independent molecules held together by an N-H...O hydrogen bond. The minor component disordered atoms of the bicyclic ring systems have been omitted for clarity. **B**, ORTEP of **11** showing two independent molecules of **11** held together by an N-H...O hydrogen bond.

ACKNOWLEDGMENTS

This research was supported by National Health and Medical Research Council (Grant APP1024314 to R.A.D), the Australian Research Council for support towards NMR and MS equipment (Grant LE0668477 and LE0237908) and financial support (Grant LP120200339 to R.A.D.). W. Loa is acknowledged for the HRESIMS measurements. F.A.E. thanks Griffith University for Ph.D. scholarships (GUPRS and GUIPRS).

A. Supplementary material

Full details of the data collection and refinement and tables of atomic coordinates, bond lengths and angles, and torsion angles have been deposited with the Cambridge Crystallographic Data Centre (CCDC 1488223 - 1488224). Copies can be obtained free of charge on application at the following address: <http://www.ccdc.cam.ac.uk/cgi-bin/catreq.cgi>. Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/>

REFERENCES

1. Newman DJ, Cragg GM. *J Nat Prod.* 2016;79:629-661.
2. Grabowski K, Baringhaus K-H, Schneider G. *Nat Prod Rep.* 2008;25:892-904.
3. Newman DJ. *J Med Chem.* 2008;51:2589-2599.
4. Ertl P, Roggo S, Schuffenhauer A. *J Chem Inf Model.* 2008;48:68-74.
5. Rosén J, Gottfries J, Muresan S, et al. *J Med Chem.* 2009;52:1953-1962.
6. Davies J. *ASM News.* 1999;65:304-310.
7. Li D, Xu S, Cai H, et al. *ChemMedChem.* 2013;8:812-818.
8. Houghton PJ. *J Ethnopharmacol.* 1988;22:121-142.
9. Yager J, Siegfried SL, DiMatteo TL. *Am J Psych.* 1999;156:1432-1438.
10. Khom S, Baburin I, Timin E, et al. *Neuropharmacology.* 2007;53:178-187.
11. Murakami N, Ye Y, Kawanishi M, et al. *Bioorg Med Chem Lett.* 2002;12:2807-2810.
12. Bos R, Hendriks H, Scheffer J, et al. *Phytomedicine.* 1998;5:219-225.
13. Hintersteiner J, Haider M, Luger D, et al. *Eur J Pharmacol.* 2014;735:123-131.
14. Circosta C, De Pasquale R, Samperi S, et al. *J Ethnopharmacol.* 2007;112:361-367.
15. Bos R, Hendriks H, Bruins A, et al. *Phytochemistry.* 1985;25:133-135.
16. Wang P-C, Hu J-M, Ran X-H, et al. *J Nat Prod.* 2009;72:1682-1685.
17. Tang Y-P, Liu X, Yu B. *J Asian Nat Prod Res.* 2003;5:257-261.
18. Tang Y, Liu X, Yu B. *J Nat Prod.* 2002;65:1949-1952.
19. Piccinelli AL, Arana S, Caceres A, et al. *J Nat Prod.* 2004;67:1135-1140.

20. Ming DS, Yang YY, He CH. *Tetrahedron Lett.* 1997;38:5205-5208.
21. Becker A, Felgentreff F, Schröder H, et al. *BMC Complement Altern Med* .2014;14:267-271.
22. Felgentreff F, Becker A, Meier B, et al. *Phytomedicine.* 2012;19:1216-1222.
23. Kopp S, Baur R, Sigel E, et al. *ChemMedChem.* 2010;5:678-681.
24. Benke D, Barberis A, Kopp S, et al. *Neuropharmacology.* 2009;56:174-181.
25. Jacobo-Herrera NJ, Vartiainen N, Bremner P, et al. *Phytother Res.* 2006;20:917-919.
26. Barnes EC, Choomuenwai V, Andrews KT, et al. *Org Biomol Chem.* 2012;10:4015-4023.
27. Choomuenwai V, Andrews KT, Davis RA. *Bioorg Med Chem.* 2012;20:7167-7174.
28. Davis RA, Carroll AR, Quinn RJ. *Aust J Chem.* 2001;54:355-359.
29. Kumar R, Sadowski MC, Levrier C, et al. *J Nat Prod.* 2015;78:914-918.
30. Davis RA, Pierens GK, Parsons PG. *Magn Reson Chem.* 2007;45:442-445.
31. Ojima I. *J Med Chem.* 2008;51:2587-2588.
32. Montalbetti CA, Falque V. *Tetrahedron.* 2005;61:10827-10852.
33. Valeur E, Bradley M. *Chem Soc Rev.* 2009;38:606-631.
34. Mishra BB, Tiwari VK. *Eur J Med Chem.* 2011;46:4769-4807.
35. ChemDraw Ultra 12.0.2.1076. www.cambridgesoft.com/software/ChemDraw; Accessed 15.02.17.
36. Lipinski CA, Lombardo F, Dominy BW, et al. *Adv Drug Deliv Rev.* 1997;23:3-25.
37. Bodanszky M. *Int J Pept Protein Res.* 1985;25:449-474.
38. Humphrey JM, Chamberlin AR. *Chem Rev.* 1997;97:2243-2266.
39. Wellenzohn B, Lessel U, Beller A, et al. *J Med Chem.* 2012;55:11031-11041.
40. Khom S, Strommer B, Ramharter J, et al. *Br J Pharmacol.* 2010;161:65-78.
41. Ramharter J, Mulzer J. *Org Lett.* 2009;11:1151-1153.
42. Birnbaum GI, Findlay JA, Krepinsky JJ. *J Org Chem.* 1978;43:272-276.
43. Nilsson N, Felding J. *Future Med Chem.* 2015;7:1853-1859.
44. Profile of Psoriasis, International Federation of Psoriasis Associations. <http://www.ifpa-pso.org>; Accessed 15.02.17.
45. Kurd SK, Gelfand JM. *J Am Acad Dermatol.* 2009;60:218-224.
46. Rothstein B, Gottlieb A. *Expert Opin Biol Ther.* 2015;16:1-10.
47. Chiricozzi A, Guttman-Yassky E, Suárez-Fariñas M, et al. *J Invest Dermatol.* 2011;131:677-687.
48. van der Bruggen T, Nijenhuis S, van Raaij E, et al. *Infect Immun.* 1999;67:3824-3829.